

Patient Information
REDEMPLO® [ree-DEM-plo]
(plozasiran)
injection, for subcutaneous use

Read this Patient Information carefully before you start taking REDEMPLO. If you have any questions about REDEMPLO, ask your healthcare provider.

What is REDEMPLO?

- REDEMPLO is an injectable prescription medicine used together with a low-fat diet to reduce triglycerides (fat in the blood) in adults with a condition that keeps the body from breaking down fats called familial chylomicronemia syndrome (FCS).
- It is not known if REDEMPLO is safe and effective in children.

Before using REDEMPLO, tell your healthcare provider about all of your medical conditions, including if you:

- are pregnant or plan to become pregnant. It is not known if REDEMPLO can harm your unborn baby. Tell your healthcare provider if you become pregnant while using REDEMPLO.
- are breastfeeding or plan to breastfeed. It is not known if REDEMPLO passes into your breast milk and if it can harm your baby. Talk with your healthcare provider about the best way to feed your baby while using REDEMPLO.

Tell your healthcare provider about all of the medicines you take including prescription and over-the-counter medicines, vitamins, and herbal supplements. Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

How should I use REDEMPLO?

- Read the **Instructions for Use** that comes with your REDEMPLO pre-filled syringe.
- Your healthcare provider will show you and/or your caregiver(s) how to inject REDEMPLO the first time.
- REDEMPLO is injected under the skin (subcutaneous use) in the front of your upper legs (thighs) or in your stomach area (abdomen). Only your healthcare provider or caregiver(s) may give you an injection in the outer area of your upper arm.
- REDEMPLO should be injected 1 time every 3 months.
- If you miss a dose, take the missing dose as soon as possible. Then inject REDEMPLO 3 months from the date of your last dose to get back on the every-3-month dosing schedule.
- Stay on a low-fat diet (less than or equal to 20 grams of fat each day) while using REDEMPLO.

What are the possible side effects of REDEMPLO?

The most common side effects of REDEMPLO include:

- high blood sugar (hyperglycemia)
- headache
- nausea
- injection site reaction (pain, redness, or swelling)

These are not all of the possible side effects of REDEMPLO. Tell your healthcare provider if you have any of the above side effects. Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or by calling 1-844-REDEMPLO (1-844-733-3675) or via <https://arrowheadpharma.com/safetyreporting>.

How should I store REDEMPLO?

- Store REDEMPLO prefilled syringe in the refrigerator between 36°F to 46°F (2°C to 8°C) in the original carton, until you are ready to use.
- REDEMPLO can also be stored at room temperature between 68°F to 77°F (20°C to 25°C) in the original carton for up to 30 days.
- **Throw away** the REDEMPLO prefilled syringe if kept at room temperature longer than 30 days.

General information about the safe and effective use of REDEMPLO.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information Leaflet. Do not use REDEMPLO for a condition for which it was not prescribed. Do not give REDEMPLO to other people, even if they have the same symptoms that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about REDEMPLO that is written for health professions.

What are the ingredients in REDEMPLO?**Active ingredients:** plozasiran sodium**Inactive ingredients:** sodium chloride, water for injection

Distributed by:

Arrowhead Pharmaceuticals, Inc.

Pasadena, CA 91105

© 2025, Arrowhead Pharmaceuticals, Inc.

All rights reserved.

This Patient Information has been approved by the U.S. Food and Drug Administration

Approved:11/2025